arXiv:1611.07137v1 [math.CO] 22 Nov 2016

On extremal multiplicative Zagreb indices of trees with given number of vertices of maximum degree

Shaohui Wang[a], Chunxiang Wang[b,*], Lin Chen[b], Jia-Bao Liu[c]

a. Department of Mathematics and Computer Science, Adelphi University, Garden City, NY 11550, USA

b. School of Mathematics and Statistics, Central China Normal University, Wuhan 430079, P.R. China

c. School of Mathematics and Physics, Anhui Jianzhu University, Hefei 230601, P.R. China

Abstract

The first multiplicative Zagreb index of a graph G is the product of the square of every vertex, while the second multiplicative Zagreb index is the product of the products of degrees of pairs of adjacent vertices. The (multiplicative) Zagreb indices are becoming the focus of considerable research in computational and analytic chemistry dating back to Gutman and Trinajstić in 1972. We explore the trees in terms of given number of vertices of maximum degree. The maximum and minimum of $\prod_1(G)$ and $\prod_2(G)$ of trees with arbitrary number of maximum degree are provided. In addition, the corresponding extremal graphs are characterized.

Keywords: Trees; Maximum Degree; Extremal bounds; Multiplicative Zagreb indices.
AMS subject classification: 05C05, 05C12

1 Introduction

Throughout this paper, we consider simple, connected and undirected graphs. Denote a graph by $G = (V, E)$, where $V = V(G)$ is called vertex set and $E = E(G)$ is called edge set. For a vertex $v \in V(G)$, the neighborhood of v is the set $N(v) = N_G(v) = \{w \in V(G), vw \in E(G)\}$, and $d_G(v)$ (or $d(v)$) denotes the degree of v with $d_G(v) = |N(v)|$. n_i is the number of vertices of degree $i \geq 0$. If a graph G contains n vertices and $n-1$ edges, then G is called a tree. For a vertex $v \in V(T)$ with $2 \leq d_T(v) \leq \Delta(T) - 1$, its edge rotating capacity is defined to be $d_T(v) - 1$. The total edge rotating capacity of a tree T is equal to the sum of the edge rotating capacities of its vertices that satisfy the condition $2 \leq d_T(v) \leq \Delta(T) - 1$. As usual, denote P_n by the path on n vertices. The maximum vertex degree in the graph G is denoted by $\Delta(G)$.

The degree sequence of G is a sequence of positive integers $\pi = (d_1, d_2, \cdots, d_n)$ if $d_i = d_G(v)(i = 1, \cdots, n)$ holds for $v \in V(G)$. In this work, we assign an order of the vertex degrees as non-increasing,

*Authors'email address: S. Wang (e-mail: shaohuiwang@yahoo.com; swang@adelphi.edu), C. Wang(e-mail: wcxiang@mailccnu.edu.cn), L. Chen(e-mail: 409137414@qq.com) J.B. Liu (e-mail: liujiabaoad@163.com).

i.e., $d_1 \geq d_2 \geq \cdots \geq d_n$. In addition, a sequence $\pi = (d_1, d_2, \cdots, d_n)$ is called a tree degree sequence if there exists a tree T such that π is its degree sequence. Furthermore, it is well known that the sequence $\pi = (d_1, d_2, \cdots, d_n)$ is a degree sequence of a tree with n vertices if and only if

$$\sum_{i=1}^{n} d_i = 2(n-1).$$

In the interdisplinary of mathemactics, chemistry and physics, molecular invariants/descriptors could be useful for the study of QSAR/QSPR and for the descriptive presentations of biological and chemical properties, such as boiling and melting points, toxicity, physico-chemical, and biological properties [1, 11, 12, 13, 20, 21]. One class of the oldest topological molecular descriptors are named as Zagreb indices [2], which are literal quantities in an expected formulas for the total π-electron energy of conjugated molecules as follows.

$$M_1(G) = \sum_{u \in V(G)} d(u)^2 \text{ and } M_2(G) = \sum_{uv \in E(G)} d(u)d(v).$$

Based on the successful considerations on these applications of Zagreb indices [3], Todeschini et al.(2010) [14, 15, 19] presented the following multiplicative variants of molecular structure descriptors:

$$\prod_1(G) = \prod_{u \in V(G)} d(u)^2 \text{ and } \prod_2(G) = \prod_{uv \in E(G)} d(u)d(v) = \prod_{u \in V(G)} d(u)^{d(u)}.$$

Recently, there are lots of articles explored multiplicative Zagreb indices in the interdisplinary of chemistry and mathematics [4, 5, 6, 7, 8, 9, 10]. Iranmanesh et al. [17] explored first and second multiplicative Zagreb indices for a class of chemical moleculor of dendrimers. Xu and Hua [16] provided a unified approach to characterize extremal maximal and minimal trees, unicyclic graphs and bicyclic graphs regarding to multiplicative Zagreb indices, respectively. Wang and Wei [19] gave the maximum and minimum indices of these indices in k-trees, and the corresponding extreme graphs are provided. Liu and Zhang [14] investigated some sharp upper bounds for $\prod_1$-index and $\prod_2$-index in terms of graph parameters such as an order, a size and a radius [18]. Kazemi [22] studied the bounds for the moments and the probability generating function of these indices in a randomly chosen molecular graph with tree structure of order n. Borovićanin et al. [23] introduced upper bounds on Zagreb indices of trees, and a lower bound for the first Zagreb index of trees with a given domination number is determined and the extremal trees are characterized as well. Borovićanin and Lampert[24] provided the maximum and minimum Zagreb indices of trees with given number of vertices of maximum degree.

Motivated by above results, in this paper we further investigate the multiplicative Zagreb indices of trees with arbitrary number of vertices of maximum degree. The maximum and minimum of $\prod_1(G)$ and $\prod_2(G)$ of trees with arbitrary number of maximum degree are provided. In addition, the corresponding extreme graphs are charaterized. Our results extends and enriches some known conclusions obtained by [24].

2 Preliminaries

It is known that each tree has at least two minimum degree vertices, named as pendent vertices, and some maximum degree vertices. It is natural to consider the trees with arbitrary number of maximum degree vertices.

Let $\mathcal{T}_{n,k}$ be the class of trees with n vertices, in which there exist k vertices having the maximum degree with $n > k > 0$. Note that the path P_n is the unique element of $\mathcal{T}_{n,n-2}$. So, in the following we consider the class $\mathcal{T}_{n,k}$ with $k \leq n-3$.

We first introduce several facts and tools, which are important in the proofs of following sections.

Proposition 2.1. *[24] If $T \in \mathcal{T}_{n,k}$ is a tree with k vertices of maximum degree Δ, then $\Delta \leq \lfloor \frac{n-2}{k} \rfloor + 1$.*

By the routine calculations, one can derive the following propositions.

Proposition 2.2. *Let $f(x) = \frac{x}{x+m}$ be a function with $m > 0$. Then $f(x)$ is increasing in $\mathbb{R}$.*

Proposition 2.3. *Let $g(x) = \frac{x^x}{(x+m)^{x+m}}$ be a function with $m > 0$. Then $g(x)$ is decreasing in $\mathbb{R}$.*

Here, by using the above algebraic tools, we are ready to provide the sharp upper and lower bounds of first multiplicative Zagreb index of such trees in section 3, and the sharp upper and lower bounds of second multiplicative Zagreb index of these trees in section 4. Some of notations and figures are used close to [24].

3 The sharp upper and lower bounds of first mutiplicative Zagreb index on the trees

In this section, we obtain the bounds of the first multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$.

3.1 The sharp upper bounds of $\prod_1$ on trees with given number of vertices of maximum degree

The first multiplicative Zagreb index of $\mathcal{T}_{n,k}$ can be routinely calculated if the degree sequence is given.

Lemma 3.1. Let T^1_{min} be a tree with minimal value of first multiplicative Zagreb index in $\mathcal{T}_{n,k}$. Then $\Delta(T^1_{min}) = \lfloor \frac{n-2}{k} \rfloor + 1$.

Proof. Let Δ be the maximum vertex degree in the tree T^1_{min}. By Proposition 2.1, we have $\Delta \leq \lfloor \frac{n-2}{k} \rfloor + 1$. Denote $\Delta_{max} = \lfloor \frac{n-2}{k} \rfloor + 1$ and $n - 2 = k\lfloor \frac{n-2}{k} \rfloor + r$, where $0 \leq r < k$.

Firstly, we assume that $\Delta < \Delta_{max}$. Denote $V(T^1_{min}) = \{v_1, \cdots, v_n\}$ by the degree sequence: $\pi = (d_1, d_2, \cdots, d_n)$. Then

$$\Delta = d_1 = \cdots = d_k = \Delta_{max} - t,\ t > 0.$$

Note that

$$\sum_{i=1}^{\Delta} n_i = n \tag{1}$$

and

$$\sum_{i=1}^{\Delta} i n_i = 2(n-1). \tag{2}$$

By the relations (1) and (2), we can obtain that

$$n_1 \geq 2 + (\Delta - 2)n_\Delta = 2 + k(\Delta - 2). \tag{3}$$

Let $n_1 = 2 + k(\Delta - 2) + n_1'$ with $n_1' \geq 0$. Using (1) we obtain

$$n = 2 + k(\Delta - 2) + n_1' + \sum_{i=2}^{\Delta-1} n_i + k,$$

which implies that

$$\sum_{i=2}^{\Delta-1} n_i + n_1' = r + kt. \tag{4}$$

Also, by the relation (2), one can calculate that

$$\sum_{i=2}^{\Delta-1} i n_i + n_1' = 2(r + kt). \tag{5}$$

By subtracting the relations (4) and (5), we have

$$\sum_{i=2}^{\Delta-1} (i-1) n_i = r + kt \geq kt \geq k. \tag{6}$$

Then the total edge rotating capacity of this tree is greater than or equal to k.

Let $v_i \in V(T_{min}^1)$ $(k < i \leq n)$ be a vertex that has positive edge rotating capacity and let d_i be its degree with $2 \leq d_i \leq \Delta - 1$. Define a tree T_1 with the vertex degree sequence $\pi_1 = (d_1^1, d_2^1, \cdots, d_n^1)$ such that $d_1^1 = \Delta + 1$, $d_i^1 = d_i - 1$ and $d_j^1 = d_j (j \in \{2, \cdots, n\}, j \neq i)$. By the definition of $\prod_1$, we have

$$\begin{aligned}
\frac{\prod_1(T_1)}{\prod_1(T_{min}^1)} &= \frac{(d_1+1)^2 d_2^2 \cdots (d_i - 1)^2 \cdots d_n^2}{d_1^2 d_2^2 \ldots d_i^2 \ldots d_n^2} \\
&= (\frac{d_1 + 1}{d_1})^2 (\frac{d_i - 1}{d_i})^2 \\
&= (\frac{\frac{(d_i-1)}{(d_i-1)+1}}{\frac{d_1}{d_1+1}})^2 \text{ (by Proposition 2.2)} \\
&< 1.
\end{aligned} \tag{7}$$

Therefore, $\prod_1(T_1) < \prod_1(T^1_{min})$. Note that T_1 has the maximum vertex degree equal to $\Delta+1$, then $T_1 \notin \mathcal{T}_{n,k}$. Since T^1_{min} has $k-1$ vertices of degree Δ, and the total positive edge rotating capacity of d_2, d_3, $\ldots$, $d_{\Delta-1}$ is at least $k-1$, by the relation (6). Then we can conclude the following statements.

We recursively proceed the above described transformations of the tree T^1_{min} for $k-1$ times on each vertex of maximum degree Δ. In every step, we could define a tree T^l with degree sequence $\pi_l = (d^l_1, \ldots, d^l_n)$ such that $d^l_l = \Delta+1$, $d^l_i = d^{l-1}_i - 1$ and $d^l_j = d^l_j - 1 (j \in \{1, \cdots, n\},\ j \neq i,\ l)$, where $l = 2, \cdots, k$ and d^{l-1}_i is the degree of any vertex $v_i \in V(T^{l-1})$ with $k < i \leq n$, which has positive edge rotating capacity (this vertex exists, because the total edge rotating capacity of the tree T^{l-1} is at least $k-l+1$). It is natural to see that after some described transformations, we obtain a tree whose degrees $d_{k+1}, \cdots, d_n$ are in an increasing order. Except for these steps, every such transformation strictly decreases the first multiplicative Zagreb index. Finally, we could get trees $T^k, T^{k-1}, \cdots, T^2 \in \mathcal{T}_{n,k}$, in which they have the maximum vertex degree equal to $\Delta+1 = (\Delta_{max}-t)+1$ and satisfy the conditions $\prod_1(T^k) < \prod_1(T^{k-1}) < \cdots < \prod_1(T^1) < \prod_1(T^1_{min})$. Considering that T^k has k vertices of degree $\Delta+1$, all of them are vertices of maximal degree.

Since $\prod_1(T^k) < \prod_1(T^1_{min})$, it contradicts the fact that T^1_{min} has the minimal first multiplicative Zagreb index in $\mathcal{T}_{n,k}$. Thus, $t=0$ and it could be considered as $\Delta = \Delta_{max} = \lfloor \frac{n-2}{k} \rfloor + 1$.

This completes the proof of $\Delta = \Delta_{max} = \lfloor \frac{n-2}{k} \rfloor + 1$. □

Based on the above proof, the following remark is immediate.

Remark 3.1. The statement of Lemma 3.1 holds for $k = n-2$, i.e., $T_{n,n-2} = P_n$.

The next theorem provides the sharp lower bound of $\prod_1(G)$ and charaterizes the exremal graphs achieving such lower bound.

Theorem 3.1. Let $T \in \mathcal{T}_{n,k}$, where $1 \leq k \leq \frac{n}{2} - 1$. Then

$$\prod_1(T) \geq \Delta^{2k}(\Delta-1)^{2p}\mu^2,$$

where the equality holds if and only if its degree sequence is $(\underbrace{\Delta, \cdots, \Delta}_{k}, \underbrace{\Delta-1, \cdots, \Delta-1}_{p}, \mu, \underbrace{1, 1, \cdots, 1}_{n-k-p-1})$ with $\Delta = \lfloor \frac{n-2}{k} \rfloor + 1$, $p = \lfloor \frac{n-2-k(\Delta-1)}{\Delta-2} \rfloor$ and $\mu = n-1-k(\Delta-1)-p(\Delta-2)$.

Proof. Let $\pi = (d_1, d_2, \cdots, d_n)$ be the vertex degree sequence of a tree T^1_{min} with minimal first multiplicative Zagreb index in $T_{n,k}$. By Lemma 3.1 we have that $d_1 = d_2 = \cdots = d_k = \Delta = \lfloor \frac{n-2}{k} \rfloor + 1$. Since $\Delta - 1 = \lfloor \frac{n-2}{k} \rfloor$, then one can obtain that $\Delta - 1$ is the integer section of $\frac{n-2}{k}$. Based on the previous lemma, let $n-2 = k(\Delta-1)+r$, where $0 \leq r < k$. From the relation (3), it follows that $n_1 \geq k(\Delta-2)+2 = n-k-r$, and n_1 has at least $n-k-r$ vertices with one degree. Therefore, $d_n = d_{n-1} = \cdots = d_{k+r+1} = 1$.

Note that $n_\Delta = k$ and $n_1 = k(\Delta+2)+2+n'_1$ with $n'_1 \geq 0$. Combining with the relations (1) and (2), and $t=0$ in the relations (5) and (6), we obtain that

$$\sum_{i=2}^{\Delta-1} n_i + n_1' = r \tag{8}$$

and

$$\sum_{i=2}^{\Delta-1} (i-1)n_i = r. \tag{9}$$

Since $n_i \geq 0$ with $i = 2, 3, \cdots, \Delta-1$, from the relation (9) it follows that $p = n_{\Delta-1} \leq \frac{r}{\Delta-2}$. Because p is a non-negative integer number, we obtain $p \leq \lfloor \frac{r}{\Delta-2} \rfloor$.

Now, suppose that $p < \lfloor \frac{r}{\Delta-2} \rfloor$ and $\lfloor \frac{r}{\Delta-2} \rfloor = t'$. Let

$$r = n_2 + 2n_3 + \cdots + (\Delta-3)n_{\Delta-2} + (\Delta-2)p = t'(\Delta-2) + y, \tag{10}$$

where $0 \leq y < (\Delta-2)$. Then $\sum_{i=2}^{\Delta-2}(i-1)n_i \geq \Delta-2$.

So there exist n_i and $n_j (2 \leq i < j \leq \Delta-2)$, where $n_i \neq 0$ and $n_j \neq 0$ or $n_i \geq 2$ (where $2 \leq i \leq \Delta-2$) and the equality (9) is satisfied. Furthermore, since $\pi = (\underbrace{\Delta, \Delta, \cdots, \Delta}_{k}, d_{k+1}, \cdots, d_{k+r}, 1, 1, \cdots, 1)$, then there exist numbers d_{k+j_1} and $d_{k+i_1} (1 \leq j_1 < i_1 \leq r)$ such that $d_{k+j_1} = j > d_{k+i_1} = i$ (or $d_{k+j_1} = d_{k+i_1} = i$, if $n_i \geq 2$).

Let $\pi' = (d_1', d_2', \cdots, d_n')$ be a sequence of positive integers such that $d'_{k+j_1} = d_{k+j_1} + 1 = j + 1$ and $d'_{k+i_1} = d_{k+i_1} + 1 = i - 1$, where $d_u' = d_u$ (for $u \neq k + j_1, u \neq k + i_1$). Clearly, $\sum_{i=1}^{n} d_i' = 2n - 2$, and π' is the vertex degree sequence of a tree T'. Also,

$$\begin{aligned} \frac{\prod_1(T')}{\prod_1(T^1_{min})} &= \frac{(j+1)^2(i-1)^2}{j^2 i^2} \\ &= (\frac{\frac{i-1}{(i-1)+1}}{\frac{j}{j+1}})^2 \text{ (by Proposition 2.3)} \\ &< 1. \end{aligned} \tag{11}$$

Therefore, $\prod_1(T') < \prod_1(T^1_{min})$. This contradicts the choice of T^1_{min} which has the minimal first multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$. Hence we conclude that $p = n_{\Delta-1} = \lfloor \frac{r}{\Delta-2} \rfloor$.

Next, by the relation (10) we obtain that $y = n_2 + 2n_3 + \cdots + (\Delta-3)n_{\Delta-2}$ (for $0 \leq y < \Delta-2$), and

$$y = r - p(\Delta-2) \leq \Delta - 3. \tag{12}$$

According to the analysis of the relation (10), it can be proved that $n_\mu = 1$ (where $2 \leq \mu \leq \Delta-2$). Then $\mu = y + 1 = r - p(\Delta-2) + 1$, i.e., $\mu = n - 1 - k(\Delta-1) - p(\Delta-2)$, and $n_i = 0$, for $i \neq \mu$, where $2 \leq i \leq \Delta-2$, since in the opposite case we can construct a tree whose $\prod_1$ is smaller than $\prod_1(T^1_{min})$ again.

Hence, the tree T^1_{min} has minimum first multiplicative Zagreb index. So we conclude that $n_\mu = 1$, and $\mu = r - p(\Delta-2) + 1$, namely, $\mu = n - 1 - k(\Delta-1) - p(\Delta-2)$.

Therefore, the tree T^1_{min} with minimum first multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$ has the vertex degree sequence $\pi = (\underbrace{\Delta, \cdots, \Delta}_{k}, \underbrace{\Delta-1, \cdots, \Delta-1}_{p}, \mu, \underbrace{1, 1, \cdots, 1}_{n-k-p-1})$. □

The first multiplicative Zagreb index of the tree $\mathcal{T}^1_{min}$ can now be routinely calculated.

Remark 3.2. The statement of Theorem 3.1 also holds for $k = n-2$, i.e., $T_{n,k} = P_n$.

3.2 The sharp lower bound of $\prod_1$ on trees with given number of vertices of maximum degree

In the following theorem we will describe the trees that have the maximal first multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$.

Lemma 3.2. Let T^1_{max} be a tree with maximal first multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$, where $1 \leq k \leq \frac{n}{2} - 1$. Then its maximum degree Δ equals to 3.

Proof. We first assume that $\Delta \geq 4$ and u is a vertex of maximum degree Δ in T^1_{max}.



Figure 1: The graphs T^1_{max} and T^1 in Lemma 3.2.

Denote $p = v_0v_1...v_{i-1}u(= v_i)v_{i+1}...v_l$ by the longest path in T^1_{max} that contains u. Also, let v_{i-1}, v_{i+1}, u_1, u_2, $...u_{\Delta-2}$ be the vertices adjacent to u in T^1_{max}, and z_1 be a pendent vertex connected to u via u_1 (it is possible that $z_1 \equiv u_1$)(Figure 1). Now we define a tree T^1 such that

$$T^1 = T^1_{max} - uu_2 + u_2z_1. \tag{13}$$

Then

$$\frac{\prod_1(T^1)}{\prod_1(T^1_{max})} = \frac{(\Delta-1)^2 2^2}{\Delta^2 1^2} = (\frac{\Delta-1}{\Delta})^2 4 > 1, \tag{14}$$

that is, $\prod_1(T^1) > \prod_1(T^1_{max})$. Obviously, the tree T^1 has $k-1$ vertices of degree Δ.

Analogously, we apply similar transformations described in the relation (13) on every vertex u of degree Δ. In each step from a tree T^i, we obtain a tree T^{i+1} with $1 \leq i \leq k-1$ that has greater first multiplicative Zagreb index than its predecessor. After k repetitions of these transformations, we arrive at the tree T^k, which has k vertices having the maximum degree $\Delta - 1$. Clearly, $T^k \in \mathcal{T}_{n,k}$ and $\prod_1(T^k) > \prod_1(T^1_{max})$.

Considering the tree maximizing $\prod_1$ in the class $\mathcal{T}_{n,k}$, it contradicts the choice of T^1_{max}. Then T^1_{max} has the maximum degree of 3, this completes our proof. □

Theorem 3.2. Let $T \in \mathcal{T}_{n,k}$ with $1 \leq k \leq \frac{n}{2} - 1$. Then

$$\prod_1(T) \leq 9^k 4^{(n-2k-2)} = (\frac{9}{16})^k 4^{n-2},$$

where the equality holds if and only if T has degree sequence $\pi = (\underbrace{3, 3, \cdots, 3}_{k}, \underbrace{2, 2, \cdots, 2}_{n-2k-2}, \underbrace{1, 1, \cdots, 1}_{k+2})$.

Proof. Let T^1_{max} be a tree with maximum $\prod_1$ in the class $\mathcal{T}_{n,k}$. By Lemma 3.2, we obtain $\Delta = 3$. So the vertex degree sequence of this tree is $\pi = (\underbrace{3, 3, \cdots, 3}_{k}, \underbrace{2, 2 \cdots, 2}_{n_2}, \underbrace{1, 1, \cdots, 1}_{n_1})$ with $k \leq \frac{n}{2} - 1$. Hence, applying to the equality (1) we obtain

$$n_1 + 2n_2 + 3k = 2(n-1) = 2(n_1 + n_2 + k) - 2. \tag{15}$$

Furthermore, since $n_3 = \Delta = k$, then

$$n_1 + 2n_2 + 3k = 2(n-1) = 2n_1 + 2n_2 + 2k - 2. \tag{16}$$

According to the relation (16), we can conclude that n_1=k+2 and n_2=n-($n_1 + n_3$)=n-2k-2. Hence,

$$\pi = (\underbrace{3, 3, \cdots, 3}_{k}, \underbrace{2, 2, \ldots, 2}_{n-2k-2}, \underbrace{1, 1, \cdots, 1}_{k+2}),$$

and

$$\prod_1(T^1_{max}) = d_1^2 d_2^2 \cdots d_n^2 = \underbrace{3^2 3^2 \cdots 3^2}_{k} \underbrace{2^2 2^2 \cdots 2^2}_{n-2k-2} \underbrace{1^2 1^2 \cdots 1^2}_{k+2} = 9^k 4^{n-2k-2} = (\frac{9}{16})^k 4^{n-2}.$$

Therefore, this completes our proof. □

Remark 3.3. Let T^k be a tree with maximal first multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$, and $T^p \in \mathcal{T}_{n,p}$, where $1 \leq k, p \leq \frac{n}{2} - 1$. If $k > p$, then $\prod_1(T^k) < \prod_1(T^p)$.

Proof. We can directly obtain by Theorem 3.2. □

4 The sharp upper and lower bounds of second mutiplicative Zagreb index on the trees

In this section, we first characterize the trees with maximal second multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$, and let $\pi = (d_1, d_2, \cdots, d_n)$ be a degree sequence of tree. To prove our main result in this section, we provide a lemma in each subsection and use it to deduce our theorems.

4.1 The sharp lower bound of $\prod_2$ on trees with given number of vertices of maximum degree

Lemma 4.1. Let T^2_{min} be a tree with minimal second multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$. Then, $\Delta(T^2_{min}) = 3$.

Proof. Suppose that $\Delta \geq 4$ and u is a vertex of maximum degree Δ in $\mathcal{T}^2_{min}$.



Figure 2: The graphs T^2_{min} and T^1 in Lemma 4.1.

Let $p = v_0v_1...v_{i-1}u(= v_i)v_{i+1}\cdots v_l$ be the longest path in T^2_{min} that contains u. Also, let v_{i-1}, v_{i+1}, u_1, u_2, $\cdots$,$u_{\Delta-2}$ be the vertices adjacent to u in T^2_{min}, and z_1 a pendent vertex connected to u via u_1 (it is possible that $z_1 \equiv u_1$)(Figure 2). Likewise, we define a tree T^1 such that

$$T^1 = T^2_{min} - uu_2 + u_2z_1. \tag{17}$$

Then

$$\begin{aligned}
\frac{\prod_2(T^1)}{\prod_2(T^2_{min})} &= \frac{(d_u-1)^{d_u-1}d_{z_1}^{d_{z_1}}}{d_u^{d_u}d_{z_1}^{d_{z_1}}} \\
&= \frac{(d_u-1)^{d_u-1}2^2}{d_u^{d_u}1^1} \\
&= 4\frac{(\Delta-1)^{\Delta-1}}{\Delta^{\Delta}} \text{(by Proposition 2.3 and } \Delta-1 \geq 3) \\
&< 4\frac{3^3}{(3+1)^{3+1}} < 1.
\end{aligned} \tag{18}$$

Therefore, $\prod_2(T^1) < \prod_2(T^2_{min})$.

In the same way, we use these transformations described in the relation (16) on every vertex v of degree Δ. Now, we transform T^2_{min} with the remaining $k-1$ vertices of the degree Δ into $\Delta-1$. After k repetitions of the transformation we arrive at the trees T^2, T^3,$\cdots$,T^k, in which k vertices have maximum degree $\Delta-1$. Obviously, $T^k \in \mathcal{T}_{n,k}$ and $\prod_2(T^k) < \prod_2(T^{k-1}) < \cdots < \prod_2(T^2_{min})$.

Thus, T^k has k the maximum vertex degree. This contradicts the choice of T^2_{min} as the tree that minimizes $\prod_2$ in the class $\mathcal{T}_{n,k}$. Therefore, $\Delta(T^2_{min}) = 3$. □

Theorem 4.1. Let $T \in \mathcal{T}_{n,k}$, where $1 \leq k \leq \frac{n}{2} - 1$. Then

$$\prod_2(T) \geq (3^3)^k (2^2)^{n-2k-2} = (\frac{27}{16})^k 4^{n-2},$$

where the equality holds if and only if T has degree sequence $\pi = (\underbrace{3, 3, \cdots, 3}_{k}, \underbrace{2, 2, \cdots, 2}_{n-2k-2}, \underbrace{1, 1, \cdots, 1}_{k+2})$.

Proof. Let T^2_{min} be a tree with minimal $\prod_2(G)$ in the class $\mathcal{T}_{n,k}$. According to Lemma 4.1, we obtain $\Delta(T^2_{min}) = 3$. Thus,

$$\pi = (\underbrace{3, 3, \cdots, 3}_{k}, \underbrace{2, 2 \cdots, 2}_{n_2}, \underbrace{1, 1, \cdots, 1}_{n_1}).$$

Clearly, $n_1 + 2n_2 + 3k = 2(n-1) = 2n_1 + 2n_2 + 2k - 2$. Then $n_1 = k+2$, $n_2 = n - n_1 - n_3 = n - 2k - 2$. Therefore, $\prod_2(T^2_{min}) = (3^3)^k (2^2)^{n_2} (1^1)^{n_1} = 27^k 4^{n-2k-2} = (\frac{27}{16})^k 4^{n-2}$. □

Remark 4.1. Let T^k be a tree with minimal second multiplicative Zagreb index in $\mathcal{T}_{n,k}$, and T^p a tree with minimal second multiplicative Zagreb index in the class $\mathcal{T}_{n,p}$, where $1 \leq k, p \leq \frac{n}{2} - 1$. If $k > p$, then $\prod_2(T^k) > \prod_2(T^p)$.

Proof. According to Theorem 4.1 we can deduce that the proposition is correct. □

4.2 The sharp upper bound of $\prod_2$ on trees with given number of vertices of maximum degree

Lemma 4.2. Let T^2_{max} be a tree with maximum second multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$, then its maximum vertex degree $\Delta(T^2_{max}) = \lfloor \frac{n-2}{k} \rfloor + 1$.

Proof. Let $\Delta = \Delta(T^2_{max})$ be the maximum vertex degree of a tree T^2_{max}. By Proposition 2.1, $\Delta \leq \lfloor \frac{n-2}{k} \rfloor + 1$. By the similar proof of Lemma 3.1, we can conclude that the tree T^1. Then,

$$\begin{aligned} \frac{\prod_2(T^1)}{\prod_2(T^2_{max})} &= \frac{(\Delta+1)^{\Delta+1}(d_i-1)^{d_i-1}}{\Delta^{\Delta} d_i^{d_i}} \\ &= \frac{\frac{(d_i-1)^{(d_i-1)}}{((d_i-1)+1)^{((d_i-1)+1)}}}{\frac{\Delta^{\Delta}}{(\Delta+1)^{\Delta+1}}} \text{(by Proposition 2.3 and } 2 \leq d_i \leq \Delta - 1 < \Delta) \\ &> 1. \end{aligned} \tag{19}$$

Therefore, $\prod_2(T^1) > \prod_2(T^2_{max})$. We can repeat remaining described transformations of the tree T^2_{max} on every of degree Δ. Therefore, we can conclude that this contradicts the fact that T^2_{max} has the maximum second multiplicative Zagreb index in the class $\mathcal{T}_{n,k}$.

This proves that $\Delta(T^2_{max}) = \lfloor \frac{n-2}{k} \rfloor + 1$. □

Theorem 4.2. Let $T \in \mathcal{T}_{n,k}$, where $1 \leq k \leq \frac{n}{2} - 1$. Then

$$\prod_2(T) \leq \Delta^{k\Delta}(\Delta-1)^{p(\Delta-1)}\mu^{\mu},$$

where the equality holds if and only if the tree T has a vertex degree sequence $\pi = (\underbrace{\Delta, \Delta, \cdots, \Delta}_{k}, \underbrace{(\Delta-1), (\Delta-1), \cdots, (\Delta-1)}_{p}, \mu, \underbrace{1, 1, \cdots, 1}_{n-k-p-1})$, for $\mu = n-1-k(\Delta-1)-p(\Delta-2)$ and $p = \lfloor \frac{n-2-k(\Delta-1)}{\Delta-2} \rfloor$.

Proof. Let T^2_{max} be a tree with maximum $\prod_2$ in the class $\mathcal{T}_{n,k}$. By Lemma 4.2, we have that $\Delta(T^2_{max}) = \lfloor \frac{n-2}{k} \rfloor + 1$. Let $p = n_{\Delta-2}$, then $p = \lfloor \frac{r}{\Delta-2} \rfloor$. Otherwise, we obtain the tree T' (using the method is analogous to the Theorem 3.1 which constructs the tree T') such that π' is the vertex degree sequence of a tree T'. Also, $\pi' = (d'_1, d'_2, \cdots, d'_n)$ is a sequence of positive integers such that $d'_{k+j_1} = d_{k+j_1}+1 = j+1$ and $d'_{k+i_1} = d_{k+i_1} + 1 = i - 1$, where $d'_u = d_u$ (for $u \neq k + j_1, u \neq k + i_1$). Thus,

$$\begin{aligned} \frac{\prod_2(T')}{\prod_2(T^2_{max})} &= \frac{(j+1)^{j+1}(i-1)^{i-1}}{j^j i^i} \\ &= \frac{\frac{(i-1)^{i-1}}{i^i}}{\frac{j^j}{(j+1)^{(j+1)}}} \text{(by Proposition 2.3)} \\ &> 1. \end{aligned} \tag{20}$$

Therefore, we obtain the tree T' such that $\prod_2$ is greater than predecessor. So $p = \lfloor \frac{r}{\Delta-2} \rfloor$. In the same way as previously, we also obtain $\prod_2(T) \leq \Delta^{k\Delta}(\Delta-1)^{p(\Delta-1)}\mu^{\mu}$.

Finally, this completes our proof. □

Acknowledgements.

The work was partially supported by the National Science Foundation of China under Grant nos. 11271149, 11371162 and 11601006, the Natural Science Foundation for the Higher Education Institutions of Anhui Province of China under Grant no. KJ2015A331. Also, it was partially supported by the Self-determined Research Funds of CCNU from the colleges basic research and operation of MOE.

References

[1] S. Klavžar, I. Gutman, The Szeged and the Wiener index of graphs, Appl. Math. Lett. 9 (1996), 45-49.

[2] I. Gutman, N. Trinajstić, Graph theory and molecular orbitals. Total -electron energy of alternant hydrocarbons, Chem. Phys. Lett. 17 (1972) 535-538.

[3] I. Gutman, On the origin of two degreeCbased topological indices, Bull. Acad. Serbe Sci. Arts. (Cl. Sci. Math.) 146 (2014) 39-52.

[4] Y. Hu, X. Li, Y. Shi, T. Xu, I. Gutman, On molecular graphs with smallest and greatest zeroth-Corder general randić index, MATCH Commun. Math. Comput. Chem. 54 (2005) 425-434.

[5] X. Li, Y. Shi, A survey on the randic index, MATCH Commun. Math. Comput. Chem 59 (2008) 127-156.

[6] Y. Shi, Note on two generalizations of the randic index, Appl. Math. Comput 265 (2015) 1019-1025.

[7] B. Furtula, I. Gutman, S. Ediz, On difference of Zagreb indices, Discrete. Appl. Math 178 (2014) 83-88.

[8] S.M. Hosamani, I. Gutman, Zagreb indices of transformation graphs and total transformation graphs, Appl. Math. Comput 247 (2014) 1156-1160.

[9] K. Xu, K.C. Das, S. Balachandran, Maximizing the Zagreb indices of (n, m)-graphs, MATCH Commun. Math. Comput. Chem 72 (2014) 641-654.

[10] J.F. Wang, F. Belardo, A lower bound for the first Zagreb index and its application, MATCH Commun. Math. Comput. Chem 74 (2015) 35-56.

[11] J. B. Liu, X. F. Pan, Asymptotic incidence energy of lattices, Physica A 422 (2015) 193-202.

[12] J. B. Liu, X. F. Pan, F. T. Hu, F. F. Hu, Asymptotic Laplacian-energy-like invariant of lattices, Appl. Math. Comput. 253 (2015) 205-214.

[13] J. B. Liu, X. F. Pan, A unified approach to the asymptotic topological indices of various lattices, Appl. Math. Comput. 270 (2015) 62-73.

[14] R. Todeschini, D. Ballabio, V. Consonni, Novel molecular descriptors based on functions of new vertex degrees, in: I. Gutman, B. Furtula (Eds.), Novel Molecular Structure Descriptors Theory and Applications I, Univ. Kragujevac, Kragujevac, (2010) 72-100.

[15] R. Todeschini, V. Consonni, New local vertex invariants and molecular descriptors based on functions of the vertex degrees, MATCH Commun. Math. Comput. Chem. 64 (2010) 359-372.

[16] K. Xu, H. Hua, A unified approach to extremal multiplicative Zagreb indices for trees, unicyclic and bicyclic graphs, MATCH Commun. Math. Comput. Chem. 68 (2012) 241-256.

[17] A. Iranmanesh, M.A. Hosseinzadeh, I. Gutman, On multiplicative Zagreb indices of graphs, Iranian J. Math. Chem. 3 (2) (2012) 145-154.

[18] J. Liu, Q. Zhang, Sharp upper bounds for multiplicative Zagreb indices, MATCH Commun. Math. Comput. Chem. 68 (2012) 231-240.

[19] S. Wang, B. Wei, Multiplicative Zagreb indices of k-trees, Discrete Appl. Math. 180 (2015) 168-175.

[20] C. Wang, S. Wang, B. Wei, Cacti with Extremal PI Index, Transactions on Combinatorics 5 (2016) 1-8.

[21] S. Wang, B. Wei, Multiplicative Zagreb indices of Cacti, Discrete Mathematics, Algorithms and Applications (2016) 1650040.

[22] R. Kazemi, Note on the multiplicative Zagreb indices, Discrete Appl. Math. 198 (2016) 147-154.

[23] B. Borovićanin, B. Furtula, On extremal Zagreb indices of trees with given domination number, Appl. Math. Comput. 279 (2016) 208-218.

[24] B. Borovićanin, T.A. Lampert, On the maximum and minimum Zagreb indices of trees with a given number of vertices of maximum degree, MATCH Commun. Math. Comput. Chem. 74 (2015) 81-96